

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number
WO 2005/077947 A1

(51) International Patent Classification⁷: **C07D 471/04**,
A61P 1/04, A61K 31/424, C07D 333/38, 333/24, 277/56,
277/30, 213/87, 213/54, 215/54, 215/12

(21) International Application Number:
PCT/EP2005/050697

(22) International Filing Date: 17 February 2005 (17.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04003606.3 18 February 2004 (18.02.2004) EP

(71) Applicant (for all designated States except US): **ALTANA
PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Kon-
stanz (DE).

(72) Inventors (for all designated States except CA, PH, US):
ZIMMERMANN, Peter, Jan; Zum Lerchental 43/1,
78315 Radolfzell (DE). **BREHM, Christof**; Richentalstr.
8, 78462 Konstanz (DE). **PALMER, Andreas**; Pirmin-
weg 1a, 78467 Konstanz (DE). **CHIESA, M., Vittoria**;
Alpenstr. 5, 78464 Konstanz (DE). **SIMON, Wolf-
gang-Alexander**; Schubertstrasse 17, 78464 Konstanz
(DE). **POSTIUS, Stefan**; Austrasse 4b, 78467 Konstanz
(DE). **KROMER, Wolfgang**; Hinterhauserstr. 5, 78464
Konstanz (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BUHR, Wilhm**
[DE/DE]; Zum Kirchenwald 7, 78465 Konstanz (DE).

(74) Agents: **WOLF, Ulrich et al.**; c/o Altana Pharma AG,
Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG)
— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 1,2,4-TRIAZOLO[4,3-A]PYRIDINES USEFUL IN THE TREATMENT OF GASTROINTESTINAL DISORDERS



(57) Abstract: The invention provides compounds of the formula (1), in which the sub-
stituents and symbols are as defined in the description. The compounds inhibit the secretion
of gastric acid.

WO 2005/077947 A1

1,2,4-TRIAZOLO[4,3-a] PYRIDINES USEFUL IN THE TREATMENT OF GASTROINTESTINAL DISORDERS.

Technical field


The invention relates to novel compounds which are used in the pharmaceutical industry as active compounds for preparing medicaments.

Prior Art

U.S. Patent 4,358,453 describes differently substituted 1,2,4-triazolo[4,3-a]pyridines, which compounds are said to be useful in the treatment of peptide ulcer.

Description of the Invention

The invention provides compounds of the formula 1



where

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group, Ar is one with R4, R5, R6 and R7 substituted mono- or bicyclic aromatic residue from the group of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, fluoro-1-4C-alkyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, fluoro-1-4C-alkyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl

R6 is hydrogen, 1-4C-alkyl or halogen and

R7 is hydrogen, 1-4C-alkyl or halogen,

wherein

aryl is phenyl or substituted phenyl with one, two or three same or different substituents from the group of 1-4C-alkyl, 1-4C-alkoxy, carboxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxy and cyano.

with the proviso that, in the case when Ar is not a 2-ethyl-6-methyl-phenyl radical, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and the salts of these compounds.

1-4C-Alkyl denotes straight-chain or branched alkyl radicals having 1 to 4 carbon atoms. Examples which may be mentioned are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

3-7C-Cycloalkyl denotes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, among which cyclopropyl, cyclobutyl and cyclopentyl are preferred.

3-7C-Cycloalkyl-1-4C-alkyl denotes one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned 3-7C-cycloalkyl radicals. Examples which may be mentioned are the cyclopropylmethyl, the cyclohexylmethyl and the cyclohexylethyl radicals.

1-4C-Alkoxy denotes radicals which, in addition to the oxygen atom, contain a straight-chain or branched alkyl radical having 1 to 4 carbon atoms. Examples which may be mentioned are the butoxy,

isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy and preferably the ethoxy and methoxy radicals.

1-4C-Alkoxy-1-4C-alkyl denotes one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the methoxymethyl, the methoxyethyl and the butoxyethyl radicals.

1-4C-Alkoxycarbonyl (-CO-1-4C-alkoxy) denotes a carbonyl group to which is attached one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the methoxycarbonyl ($\text{CH}_3\text{O}-\text{C}(\text{O})-$) and the ethoxycarbonyl ($\text{CH}_3\text{CH}_2\text{O}-\text{C}(\text{O})-$) radicals.

2-4C-Alkenyl denotes straight-chain or branched alkenyl radicals having 2 to 4 carbon atoms. Examples which may be mentioned are the 2-butenyl, 3-butenyl, 1-propenyl and the 2-propenyl (allyl) radicals.

2-4C-Alkynyl denotes straight-chain or branched alkynyl radicals having 2 to 4 carbon atoms. Examples which may be mentioned are the 2-butyne, the 3-butyne and, preferably, the 2-propyne (propargyl radicals).

Fluoro-1-4C-alkyl denotes one of the abovementioned 1-4C-alkyl radicals which is substituted by one or more fluorine atoms. An example which may be mentioned is the trifluoromethyl radical.

Hydroxy-1-4C-alkyl denotes abovementioned 1-4C-alkyl radicals which are substituted by a hydroxyl group. Examples which may be mentioned are the hydroxymethyl, the 2-hydroxyethyl and the 3-hydroxypropyl radicals.

For the purpose of the invention, halogen is bromine, chlorine and fluorine.

1-4C-Alkoxy-1-4C-alkoxy denotes one of the abovementioned 1-4C-alkoxy radicals which is substituted by a further 1-4C-alkoxy radical. Examples which may be mentioned are the radicals 2-(methoxy)ethoxy ($\text{CH}_3-\text{O}-\text{CH}_2-\text{CH}_2-\text{O}-$) and 2-(ethoxy)ethoxy ($\text{CH}_3-\text{CH}_2-\text{O}-\text{CH}_2-\text{CH}_2-\text{O}-$).

1-4C-Alkoxy-1-4C-alkoxy-1-4C-alkyl denotes one of the abovementioned 1-4C-alkoxy-1-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy radicals. An example which may be mentioned is the radical 2-(methoxy)ethoxymethyl ($\text{CH}_3-\text{O}-\text{CH}_2-\text{CH}_2-\text{O}-\text{CH}_2-$).

Fluoro-1-4C-alkoxy-1-4C-alkyl denotes one of the abovementioned 1-4C-alkyl radicals which is substituted by a fluoro-1-4C-alkoxy radical. Here, fluoro-1-4C-alkoxy denotes one of the abovementioned 1-4C-alkoxy radicals which is fully or predominantly substituted by fluorine. Examples of fully or predominantly fluorine-substituted 1-4C-alkoxy which may be mentioned are the 1,1,1,3,3,3-hexafluoro-2-propoxy, the 2-trifluoromethyl-2-propoxy, the 1,1,1-trifluoro-2-propoxy, the perfluoro-tert-

butoxy, the 2,2,3,3,4,4,4-heptafluoro-1-butoxy, the 4,4,4-trifluoro-1-butoxy, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,2,2-trifluoroethoxy, in particular the 1,1,2,2-tetrafluoroethoxy, the 2,2,2-trifluoroethoxy, the trifluoromethoxy and preferably the difluoromethoxy radicals.

1-7C-Alkyl denotes straight-chain or branched alkyl radicals having 1 to 7 carbon atoms. Examples which may be mentioned are the heptyl, isoheptyl-(5-methylhexyl), hexyl, isoheptyl-(4-methylpentyl), neoheptyl-(3,3-dimethylbutyl), pentyl, isopentyl-(3-methylbutyl), neopentyl-(2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and methyl radicals.

1-4C-Alkylcarbonyl denotes a radical which, in addition to the carbonyl group, contains one of the abovementioned 1-4C-alkyl radicals. An example which may be mentioned is the acetyl radical.

2-4C-Alkenyloxy denotes a radical which, in addition to the oxygen atom, contains a 2-4C-alkenyl radical. An example which may be mentioned is the allyloxy radical.

Carboxy-1-4C-alkyl denotes, for example, the carboxymethyl ($-\text{CH}_2\text{COOH}$) or the carboxyethyl ($-\text{CH}_2\text{CH}_2\text{COOH}$) radical.

1-4C-Alkoxy-1-4C-alkyl denotes one of the abovementioned 1-4C-alkyl radicals which is substituted by one of the abovementioned 1-4C-alkoxy radicals. An example which may be mentioned is the ethoxycarbonylmethyl ($\text{CH}_3\text{CH}_2\text{OC(O)CH}_2-$) radical.

Aryl-1-4C-alkyl denotes an aryl-substituted 1-4C-alkyl radical. An example which may be mentioned is the benzyl radical.

Aryl-1-4C-alkoxy denotes an aryl-substituted 1-4C-alkoxy radical. An example which may be mentioned is the benzyloxy radical.

Mono- or di-1-4C-alkylamino radicals contain, in addition to the nitrogen atom, one or two of the abovementioned 1-4C-alkyl radicals. Preference is given to di-1-4C-alkylamino and in particular to dimethyl-, diethyl- or diisopropylamino.

1-4C-Alkylcarbonylamino denotes an amino group to which a 1-4C-alkylcarbonyl radical is attached. Examples which may be mentioned are the propionylamino ($\text{C}_3\text{H}_7\text{C(O)NH-}$) and the acetylamino (acetamido, $\text{CH}_3\text{C(O)NH-}$) radicals.

1-4C-Alkoxy-1-4C-alkylamino denotes an amino radical which is substituted by one of the abovementioned 1-4C-alkoxy radicals. Examples which may be mentioned are the ethoxycarbonylamino and the methoxycarbonylamino radicals.

1-4C-Alkoxy-1-4C-alkoxycarbonyl denotes a carbonyl group to which one of the abovementioned 1-4C-alkoxy-1-4C-alkoxy radicals is attached. Examples which may be mentioned are the 2-(methoxy)ethoxycarbonyl ($\text{CH}_3\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$) and the 2-(ethoxy)ethoxycarbonyl ($\text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$) radicals.

1-4C-Alkoxy-1-4C-alkoxycarbonylamino denotes an amino radical which is substituted by one of the abovementioned 1-4C-alkoxy-1-4C-alkoxycarbonyl radicals. Examples which may be mentioned are the 2-(methoxy)ethoxycarbonylamino and the 2-(ethoxy)ethoxycarbonylamino radicals.

Radicals Ar which may be mentioned are, for example, the following substituents: 4-acetoxyphenyl, 4-acetamidophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 3-benzyloxy-4-methoxyphenyl, 4-benzyloxy-3-methoxyphenyl, 3,5-bis(trifluoromethyl)phenyl, 4-butoxyphenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-chloro-6-fluorophenyl, 3-chloro-4-fluorophenyl, 2-chloro-5-nitrophenyl, 4-chloro-3-nitrophenyl, 3-(4-chlorophenoxy)phenyl, 2,4-dichlorophenyl, 3,4-difluorophenyl, 2,4-dihydroxyphenyl, 2,6-dimethoxyphenyl, 3,4-dimethoxy-5-hydroxyphenyl, 2,5-dimethylphenyl, 3-ethoxy-4-hydroxyphenyl, 2-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-hydroxy-5-nitrophenyl, 3-methoxy-2-nitrophenyl, 3-nitrophenyl, 2,3,5-trichlorophenyl, 2,4,6-trihydroxyphenyl, 2,3,4-trimethoxyphenyl, 2-hydroxy-1-naphthyl, 2-methoxy-1-naphthyl, 4-methoxy-1-naphthyl, 1-methyl-2-pyrrolyl, 2-pyrrolyl, 3-methyl-2-pyrrolyl, 3,4-dimethyl-2-pyrrolyl, 4-(2-methoxycarbonyl)ethyl-3-methyl-2-pyrrolyl, 5-ethoxycarbonyl-2,4-dimethyl-3-pyrrolyl, 3,4-dibromo-5-methyl-2-pyrrolyl, 2,5-dimethyl-1-phenyl-3-pyrrolyl, 5-carboxy-3-ethyl-4-methyl-2-pyrrolyl, 3,5-dimethyl-2-pyrrolyl, 2,5-dimethyl-1-(4-trifluoromethylphenyl)-3-pyrrolyl, 1-(2,6-dichloro-4-trifluoromethylphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-pyrrolyl, 1-(2-fluorophenyl)-2-pyrrolyl, 1-(4-trifluoromethoxyphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-pyrrolyl, 1-(4-ethoxycarbonyl)-2,5-dimethyl-3-pyrrolyl, 5-chloro-1,3-dimethyl-4-pyrazolyl, 5-chloro-1-methyl-3-trifluoromethyl-4-pyrazolyl, 1-(4-chlorobenzyl)-5-pyrazolyl, 1,3-dimethyl-5-(4-chlorophenoxy)-4-pyrazolyl, 1-methyl-3-trifluoromethyl-5-(3-trifluoromethylphenoxy)-4-pyrazolyl, 4-methoxycarbonyl-1-(2,6-dichlorophenyl)-5-pyrazolyl, 5-allyloxy-1-methyl-3-trifluoromethyl-4-pyrazolyl, 5-chloro-1-phenyl-3-trifluoromethyl-4-pyrazolyl, 3,5-dimethyl-1-phenyl-4-imidazolyl, 4-bromo-1-methyl-5-imidazolyl, 2-butylimidazolyl, 1-phenyl-1,2,3-triazol-4-yl, 3-indolyl, 4-indolyl, 7-indolyl, 5-methoxy-3-indolyl, 5-benzyloxy-3-indolyl, 1-benzyl-3-indolyl, 2-(4-chlorophenyl)-3-indolyl, 7-benzyloxy-3-indolyl, 6-benzyloxy-3-indolyl, 2-methyl-5-nitro-3-indolyl, 4,5,6,7-tetrafluoro-3-indolyl, 1-(3,5-difluorobenzyl)-3-indolyl, 1-methyl-2-(4-trifluorophenoxy)-3-indolyl, 1-methyl-2-benzimidazolyl, 5-nitro-2-furyl, 5-hydroxymethyl-2-furyl, 2-furyl, 3-furyl, 5-(2-nitro-4-trifluoromethylphenyl)-2-furyl, 4-ethoxycarbonyl-5-methyl-2-furyl, 5-(2-trifluoromethoxyphenyl)-2-furyl, 5-(4-methoxy-2-nitrophenyl)-2-furyl, 4-bromo-2-furyl, 5-dimethylamino-2-furyl, 5-bromo-2-furyl, 5-sulfo-2-furyl, 2-benzofuryl, 2-thienyl, 3-thienyl, 3-methyl-2-thienyl, 4-bromo-2-thienyl, 5-bromo-2-thienyl, 5-nitro-2-thienyl, 5-methyl-2-thienyl, 5-(4-methoxyphenyl)-2-thienyl, 4-methyl-2-thienyl, 3-phenoxy-2-thienyl, 5-carboxy-2-thienyl, 2,5-dichloro-3-thienyl, 3-methoxy-2-thienyl, 2-benzothienyl, 3-methyl-2-benzothienyl, 2-bromo-5-chloro-3-benzothienyl, 2-thiazolyl, 2-amino-4-chloro-5-thiazolyl, 2,4-dichloro-5-thiazolyl, 2-diethylamino-5-thiazolyl, 3-methyl-4-nitro-5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 6-methyl-2-pyridyl, 3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridyl, 2,6-dichloro-4-pyridyl, 3-chloro-5-

trifluoromethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 4-(4-chlorophenyl)-3-pyridyl, 2-chloro-5-methoxy-carbonyl-6-methyl-4-phenyl-3-pyridyl, 2-chloro-3-pyridyl, 6-(3-trifluoromethylphenoxy)-3-pyridyl, 2-(4-chlorophenoxy)-3-pyridyl, 2,4-dimethoxy-5-pyrimidine, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 2-chloro-3-quinolinyl, 2-chloro-6-methoxy-3-quinolinyl, 8-hydroxy-2-quinolinyl and 4-isoquinolinyl.

Suitable salts of compounds of the formula 1 are – depending on the substitution – in particular all acid addition salts. Particular mention may be made of the pharmacologically acceptable salts of the inorganic and organic acids customarily used in pharmacy. Those suitable are water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are employed in the salt preparation in an equimolar ratio or in a ratio differing therefrom, depending on whether the acid is a mono- or polybasic acid and on which salt is desired.

Pharmacologically unacceptable salts, which can be initially obtained, for example, as process products in the preparation of the compounds according to the invention on an industrial scale, are converted into pharmacologically acceptable salts by processes known to the person skilled in the art.

It is known to the person skilled in the art that the compounds according to the invention and their salts can, for example when they are isolated in crystalline form, comprise varying amounts of solvents. The invention therefore also embraces all solvates and, in particular, all hydrates of the compounds of the formula 1, and all solvates and, in particular, all hydrates of the salts of the compounds of the formula 1.

Compounds which are to be emphasized are those of the formula 1, where

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

Ar is one with R4, R5, R6 and R7 substituted mono- or bicyclic aromatic residue from the group of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

R6 is hydrogen,

R7 is hydrogen,

with the proviso that, in the case when Ar is not a 2-ethyl-6-methyl-phenyl radical, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and the salts of these compounds.

Particular mention may be made of those compounds of the formula 1,

where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical -Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted monocyclic aromatic residue selected from the group of phenyl, pyridinyl, thiophenyl, furanyl and pyrrolyl,

wherein

R4 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

R5 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

R6 is hydrogen

R7 is hydrogen

and the salts of these compounds.

Particular mention may also be made of those compounds of the formula 1,

where

- R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,
 where
 Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical
 R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 Ar is one with R4, R5, R6 and R7 substituted phenyl,
 wherein
 R4 is hydrogen or 1-4C-alkyl
 R5 is hydrogen or 1-4C-alkyl
 R6 is hydrogen
 R7 is hydrogen
 and the salts of these compounds.

Particular emphasis is given to compounds of the formula 1,
 where

- R1 1-4C-alkyl
 R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,
 where
 Res is a imidazo or morpholino radical and is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical
 R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 Ar is one with R4, R5, R6 and R7 substituted phenyl,
 wherein
 R4 is hydrogen or 1-4C-alkyl
 R5 is hydrogen or 1-4C-alkyl
 R6 is hydrogen
 R7 is hydrogen
 and the salts of these compounds.

Particular emphasis is also given to compounds of the formula 1,
 where

- R1 1-4C-alkyl
 R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR31R32,

where

R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted phenyl,

wherein

R4 is hydrogen or 1-4C-alkyl


R5 is hydrogen or 1-4C-alkyl

R6 is hydrogen

R7 is hydrogen

and the salts of these compounds.

Among the compounds of the formula 1, those compounds of the formula 1-a are preferred.



Compounds of the formula 1-a which are to be emphasized are those,

in which

R1^{1b} is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl
 R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl
 with the proviso that, in the case when R4 is not ethyl and R5 is not methyl, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and the salts of these compounds.

Particular mention may be made of those compounds of the formula 1-a,

where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R4 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

R5 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

and the salts of these compounds.

Particular mention may also be made of those compounds of the formula 1-a,

where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl; hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R4 is hydrogen or 1-4C-alkyl

R5 is hydrogen or 1-4C-alkyl

and the salts of these compounds.

Compounds of the formula 1-a which are to be particularly emphasized are those,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR₃₁R₃₂,

where

Res is a imidazo or morpholino radical and is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R₃₁ is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R₃₂ is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R4 is hydrogen or 1-4C-alkyl

R5 is hydrogen or 1-4C-alkyl

and the salts of these compounds.

Compounds of the formula 1-a which are also to be particularly emphasized are those,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR₃₁R₃₂,

where

R₃₁ is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,


R₃₂ is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R4 is hydrogen or 1-4C-alkyl

R5 is hydrogen or 1-4C-alkyl

and the salts of these compounds.

Among the compounds of the formula 1-a, those compounds of the formula 1-b are particularly preferred



Compounds of the formula 1-b which are to be emphasized are those,

in which

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR₃₁R₃₂,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

and the salts of these compounds.

Particular mention may be made of those compounds of the formula 1-b,

where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

and the salts of these compounds.

Compounds of the formula 1-b which are to be emphasized are those,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo or morpholino radical and is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

and the salts of these compounds.

Compounds of the formula 1-b which are to be particularly emphasized are those, where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR31R32,

where


R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

and the salts of these compounds.

The compounds of the formula 1 according to the invention can be prepared as described in an exemplary manner in the examples below, or starting from appropriate starting materials using analogous process steps or as illustrated quite generally in the scheme 1 below. Compounds of the formula 2 can be transformed to compounds of the formula 3 in a manner known per se to the person skilled in the art using standard reaction conditions, like for example using hydrogen / Pd(0). The arylation of compounds of the formula 3 to compounds of the formula 1 is carried out in manner known to the person skilled in the art using a suitable Ar-CH₂-X reactant carrying a suitable leaving group X, like for example a chlorine atom.

Scheme 1:




The derivatization, if any, of the compounds obtained according to the above Scheme 1 (e.g. conversion of a group R2 into another group R2) is likewise carried out in a manner known to the expert. For example, if compounds of the formula 1 where R2 = carboxyl or -CO-NR31R32 are desired, an appropriate derivatization can be performed in a manner known to the expert (for example conversion of an ester into a carboxylic acid and further transformation into an amide) at the stage of the compounds of formula 2 or 3 or more conveniently at a later point in time, for example conversion of a compound of the formula 1 into another compound of the formula 1.

If compounds of the formula 1 where R2 = hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl or the radical Res-1-4C-alkyl are desired, an appropriate derivatization can be performed in a manner known to the expert (for example conversion of an ester into an alcohol followed by chlorination of the alcohol and any desired substitution of the chlorine atom, like for example an etherification to form compounds of the

formula 1 with R2 = 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl or a nucleophilic substitution to form compounds of the formula 1 with R2 = amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl or the radical Res-1-4C-alkyl).


Compounds of the formula 2 can be obtained in a manner known per se to the person skilled in the art, for example in analogy to the synthesis described in J. Org. Chem., 1966, 31, 251, or J. Heterocycl. Chem., 1970, 7, 1019, by cyclization of compounds of the formula 4 under suitable conditions in the presence of a suitable ortho-ester carrying suitable substituents Z, like for example methyl groups (scheme 2).

Scheme 2



Compounds of the formula 4 are known, for example from J. Heterocycl. Chem., 1970, 7, 1019, or can be prepared in an analogous manner by reactions known per se to the person skilled in the art or in a manner as shown in a general way in scheme 3.

Scheme 3



The examples below serve to illustrate the invention in more detail without limiting it. Further compounds of the formula 1, whose preparation is not described explicitly, can likewise be prepared in an analogous manner or in a manner known per se to the person skilled in the art, using customary process techniques. The compounds named expressly as examples, and the salts of these compounds, are preferred subject matter of the invention. The abbreviation min stands for minute(s), h stands for hour(s) and m.p. stands for melting point.

Examples**I. Final Products****1. 8-(2-Ethyl-6-methyl-benzylamino)-3-methyl[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid methyl ester**

To a solution of 3.80 g (18.4 mmol) 3-methyl-8-amino[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid methyl ester in DMF (50 ml) is added 3.73 g (mmol) 2-ethyl-6-methylbenzylchloride and then portionwise 0.88 g (22.1 mmol) sodium hydride (60 % dispersion in mineral oil). This mixture is stirred for 1 h at ambient temperature. Afterwards the reaction is quenched by pouring it into a saturated aqueous ammonium chloride solution. This mixture is extracted three times with ethyl acetate. The combined organic layers are concentrated in vacuo and the crude product is purified by column chromatography (dichloromethane / methanol: 100 / 3) to give 2.60 g (7.68 mmol / 42 %) of the title compound with a melting point of 214.4°C (ethyl acetate).

2. 8-(2-Ethyl-6-methyl-benzylamino)-3-methyl[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid

To a suspension of 2.50 g (7.40 mmol) 8-(2-ethyl-6-methyl-benzylamino)-3-methyl[1,2,4]triazolo-[4,3-a]pyridine-6-carboxylic acid methyl ester in dioxane (25 ml) are added 3.75 ml (7.50 mmol) of a sodium hydroxide solution (2N) and the mixture is stirred at 80°C for 5 h. Subsequently the reaction is quenched by pouring the reaction mixture into an ice cooled, saturated aqueous ammonium chloride solution. This mixture is extracted two times with dichloromethane and methanol (13 / 1). The combined organic layers are concentrated in vacuo and purified by column chromatograph (dichloromethane / methanol: 13 / 1 to 3 / 1) to give 2.40 g (7.40 mmol / 99 %) of the title compound with a melting point of 328.1°C (dichloromethane / methanol).

3. 8-(2-Ethyl-6-methylbenzylamino)-N,N,3-trimethyl[1,2,4]triazolo[4,3-a]pyridine-6-carboxamide

To a suspension of 1.00 g (3.08 mmol) 8-(2-ethyl-6-methyl-benzylamino)-3-methyl[1,2,4]triazolo-[4,3-a]pyridine-6-carboxylic acid in THF and DMF (5 / 1: 12 ml) is added 1.14 g (6.80 mmol) N,N-carbonyldiimidazole. This mixture is stirred at 25°C for 1 h and at 50°C for further 30 min. After cooling to ambient temperature, 40 mmol dimethylamine (20 ml of a 2M solution in THF) are added to the reaction mixture which is then stirred at 25°C for further 72 h. Subsequently the reaction mixture is concentrated in vacuo and the crude product is purified by column chromatography (dichloromethane / methanol: 100 / 3) to give 0.77 g (2.19 mmol / 71 %) of the title compound with a melting point of 201.3°C (diethyl ether).

4. 8-(2-Ethyl-6-methylbenzylamino)-N-(2-methoxyethyl)-3-methyl[1,2,4]triazolo-[4,3-a]pyridine-6-carboxamide

To a suspension of 1.00 g (3.08 mmol) 8-(2-ethyl-6-methyl-benzylamino)-3-methyl[1,2,4]triazolo-[4,3-a]pyridine-6-carboxylic acid in THF and DMF (5 / 1: 18 ml) is added 1.14 g (6.80 mmol) N,N-carbonyldiimidazole. This mixture is stirred at 25°C for 1 h, at 50°C for 30 min. and again at 25°C for further 2 h. Afterwards, 2.00 ml (23.0 mmol) 2-methoxyethylamine are added to the reaction mixture which is then stirred at 25°C for 18 h. Subsequently the reaction mixture is concentrated in vacuo and the crude product is purified by column chromatography (dichloromethane / methanol: 100 / 3) to yield 0.80 g (2.10 mmol / 68 %) of the title compound with a melting point of 132.0°C (diethyl ether).

II. Starting compounds and Intermediates

A. 6-Chloro-5-nitro-nicotinic acid methyl ester

To a suspension of 49.0 g (0.27 mol) 6-hydroxy-5-nitro-nicotinic acid in thionyl chloride (240 ml) are added 2 ml of DMF. This mixture is stirred at 60°C until the evolution of gas has ended. Then it is stirred at 80°C for further 18 h. Residual thionyl chloride is removed under vacuo and the resulting residue is coevaporated three times with toluene. Subsequently this reaction mixture is dissolved in dichloromethane (100 ml) and cooled to 0°C before methanol (55.5 ml) is dropwise added. The precipitated solid is filtered off and dried under vacuo at 50°C to give 27.6 g (13.7 mmol / 52 %) of the title compound as a light yellow solid with a melting point of 78°C (dichloromethane / methanol).

B. 6-Hydrazino-5-nitro-nicotinic acid methyl ester

To a at 15°C cooled solution of 30.0 g (0.14 mol) 6-chloro-5-nitro-nicotinic acid methyl ester in dioxane (600ml) is added hydrazine hydrate (21.5 ml). During the addition the reaction mixture is warmed up to 25°C and is stirred for further 3 h. The reaction is quenched by pouring the reaction mixture into a saturated aqueous ammonium chloride solution. The precipitated solid is filtered off and dried under vacuo at 50°C to give 26.5 g (0.12 mol / 90 %) of the title compound as a red solid.

¹H-NMR (200MHz, d⁶-DMSO): δ = 3.85 (s, 3 H), 8.69 (d, 1 H), 8.90 (d, 1 H).

C. 3-Methyl-8-nitro[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid methyl ester

A suspension of 14.4 g (67.9 mmol) 6-hydrazino-5-nitro-nicotinic acid methyl ester in trimethyl orthoacetate (280 ml) is stirred at 78°C for 18 h and at 25°C for further 25 h. The precipitated crude product is filtered off and purified by chromatography (dichloromethane / methanol: 100 / 3 to 13 / 1) to give 6.37 g (26.9 mmol / 40 %) of the title product as a yellow solid.

¹H-NMR (200MHz, CDCl₃): δ = 2.91 (s, 3 H), 4.06 (s, 3 H), 8.78 (d, 1 H), 8.93 (d, 1 H).

D. 3-Methyl-8-amino[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid methyl ester

A suspension of 10.0 g (42.3 mmol) 3-methyl-8-nitro[1,2,4]triazolo[4,3-a]pyridine-6-carboxylic acid methyl ester, 10 g (9.40 mmol) palladium on carbon (10 % on carbon) and 44.0 ml (0.46 mol) cyclohexadiene in a mixture of ethanol and ethyl acetate (1 / 1: 600ml) is stirred at 60°C for 4 h. Subsequently the catalyst is filtered off and the reaction mixture is concentrated in vacuo. The crude product is suspended in acetone and filtered off to give 5.00 g (24.3 mmol / 57 %) of the title product.

¹H-NMR (200MHz, CDCl₃): δ = 2.78 (s, 3 H), 3.95 (s, 3 H), 6.98 (s, 1 H), 8.11 (s, 1 H).

Commercial utility

The compounds of the formula 1 and their salts have valuable pharmacological properties which make them commercially utilizable. In particular, they exhibit marked inhibition of gastric acid secretion and an excellent gastric and intestinal protective action in warm-blooded animals, in particular humans. In this connection, the compounds according to the invention are distinguished by a high selectivity of action, an advantageous duration of action, a particularly good enteral activity, the absence of significant side effects and a large therapeutic range.

"Gastric and intestinal protection" in this connection is understood as meaning the prevention and treatment of gastrointestinal diseases, in particular of gastrointestinal inflammatory diseases and lesions (such as, for example, gastric ulcer, peptic ulcer, including peptic ulcer bleeding, duodenal ulcer, gastritis, hyperacidic or medicament-related functional dyspepsia), which can be caused, for example, by microorganisms (e.g. *Helicobacter pylori*), bacterial toxins, medicaments (e.g. certain antiinflammatories and antirheumatics, such as NSAIDs and COX-inhibitors), chemicals (e.g. ethanol), gastric acid or stress situations. "Gastric and intestinal protection" is understood to include, according to general knowledge, gastroesophageal reflux disease (GERD), the symptoms of which include, but are not limited to, heartburn and/or acid regurgitation.

In their excellent properties, the compounds according to the invention surprisingly prove to be clearly superior to the compounds known from the prior art in various models in which the antiulcerogenic and the antisecretory properties are determined. On account of these properties, the compounds of the formula 1 and their pharmacologically acceptable salts are outstandingly suitable for use in human and veterinary medicine, where they are used, in particular, for the treatment and/or prophylaxis of disorders of the stomach and/or intestine.

A further subject of the invention are therefore the compounds according to the invention for use in the treatment and/or prophylaxis of the abovementioned diseases.

The invention likewise includes the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the abovementioned diseases.

The invention furthermore includes the use of the compounds according to the invention for the treatment and/or prophylaxis of the abovementioned diseases.

A further subject of the invention are medicaments which comprise one or more compounds of the formula 1 and/or their pharmacologically acceptable salts.

The medicaments are prepared by processes which are known per se and familiar to the person skilled in the art. As medicaments, the pharmacologically active compounds according to the invention (=

active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries or excipients in the form of tablets, coated tablets, capsules, suppositories, patches (e.g. as TTS), emulsions, suspensions or solutions, the active compound content advantageously being between 0.1 and 95% and it being possible to obtain a pharmaceutical administration form exactly adapted to the active compound and/or to the desired onset and/or duration of action (e.g. a sustained-release form or an enteric form) by means of the appropriate selection of the auxiliaries and excipients.

The auxiliaries and excipients which are suitable for the desired pharmaceutical formulations are known to the person skilled in the art on the basis of his/her expert knowledge. In addition to solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound excipients, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, colorants or, in particular, permeation promoters and complexing agents (e.g. cyclodextrins).

The active compounds can be administered orally, parenterally or percutaneously.

In general, it has proven advantageous in human medicine to administer the active compound(s) in the case of oral administration in a daily dose of approximately 0.01 to approximately 20, preferably 0.05 to 5, in particular 0.1 to 1.5, mg/kg of body weight, if appropriate in the form of several, preferably 1 to 4, individual doses to achieve the desired result. In the case of a parenteral treatment, similar or (in particular in the case of the intravenous administration of the active compounds), as a rule, lower doses can be used. The establishment of the optimal dose and manner of administration of the active compounds necessary in each case can easily be carried out by any person skilled in the art on the basis of his/her expert knowledge.

If the compounds according to the invention and/or their salts are to be used for the treatment of the abovementioned diseases, the pharmaceutical preparations can also contain one or more pharmacologically active constituents of other groups of medicaments, for example: tranquillizers (for example from the group of the benzodiazepines, for example diazepam), spasmolytics (for example, biefamiverine or camylofine), anticholinergics (for example, oxyphenyclimine or phencarbamide), local anesthetics, (for example, tetracaine or procaine), and, if appropriate, also enzymes, vitamins or amino acids.

To be emphasized in this connection is in particular the combination of the compounds according to the invention with pharmaceuticals which inhibit acid secretion, such as, for example, H_2 blockers (e.g. cimetidine, ranitidine), H^+/K^+ ATPase inhibitors (e.g. omeprazole, pantoprazole), or further with so-called peripheral anticholinergics (e.g. pirenzepine, telenzepine) and with gastrin antagonists with the aim of increasing the principal action in an additive or super-additive sense and/or of eliminating or of decreasing the side effects, or further the combination with antibacterially active substances (such as, for example, cephalosporins, tetracyclines, penicillins, macrolides, nitroimidazoles or alternatively

bismuth salts) for the control of *Helicobacter pylori*. Suitable antibacterial co-components which may be mentioned are, for example, mezlocillin, ampicillin, amoxicillin, cefalothin, cefoxitin, cefotaxime, imipenem, gentamycin, amikacin, erythromycin, ciprofloxacin, metronidazole, clarithromycin, azithromycin and combinations thereof (for example clarithromycin + metronidazole).

In view of their excellent gastric and intestinal protection action, the compounds of formula 1 are suited for a free or fixed combination with those medicaments (e.g. certain antiinflammatories and antirheumatics, such as NSAIDs), which are known to have a certain ulcerogenic potency. In addition, the compounds of formula 1 are suited for a free or fixed combination with motility-modifying drugs.

Pharmacology

The excellent gastric protective action and the gastric acid secretion-inhibiting action of the compounds according to the invention can be demonstrated in investigations on animal experimental models. The compounds according to the invention investigated in the model mentioned below have been provided with numbers which correspond to the numbers of these compounds in the examples.

Testing of the secretion-inhibiting action on the perfused rat stomach

In Table A which follows, the influence of the compounds according to the invention on the pentagastrin-stimulated acid secretion of the perfused rat stomach after intraduodenal administration in vivo is shown.

Table A:

No.	Dose ($\mu\text{mol/kg}$) i.d.	Inhibition of acid secretion (%)
3	1.0	> 30
4	1.0	> 30

Methodology

The abdomen of anesthetized rats (CD rat, female, 200-250 g; 1.5 g/kg i.m. urethane) was opened after tracheotomy by a median upper abdominal incision and a PVC catheter was fixed transorally in the esophagus and another via the pylorus such that the ends of the tubes just projected into the gastric lumen. The catheter leading from the pylorus led outward into the right abdominal wall through a side opening.


After thorough rinsing (about 50-100 ml), warm (37°C) physiological NaCl solution was continuously passed through the stomach (0.5 ml/min, pH 6.8-6.9; Braun-Unita I). The pH (pH meter 632, glass electrode EA 147; $\phi = 5$ mm, Metrohm) and, by titration with a freshly prepared 0.01N NaOH solution to pH 7 (Dosimat 665 Metrohm), the secreted HCl were determined in the effluent in each case collected at an interval of 15 minutes.

The gastric secretion was stimulated by continuous infusion of 1 $\mu\text{g/kg}$ (= 1.65 ml/h) of i.v. pentagastrin (left femoral vein) about 30 min after the end of the operation (i.e. after determination of 2 preliminary fractions). The substances to be tested were administered intraduodenally in a 2.5 ml/kg liquid volume 60 min after the start of the continuous pentagastrin infusion.

The body temperature of the animals was kept at a constant 37.8-38°C by infrared irradiation and heat pads (automatic, stepless control by means of a rectal temperature sensor).

Claims

1. A compound of the formula 1



where

- R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

- Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

- R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

- R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

- R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

- R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

- Ar is one with R4, R5, R6 and R7 substituted mono- or bicyclic aromatic residue from the group of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

- R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, fluoro-1-4C-alkyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, fluoro-1-4C-alkyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl

R6 is hydrogen, 1-4C-alkyl or halogen and

R7 is hydrogen, 1-4C-alkyl or halogen,

wherein

aryl is phenyl or substituted phenyl with one, two or three same or different substituents from the group of 1-4C-alkyl, 1-4C-alkoxy, carboxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxy and cyano.

with the proviso that, in the case when Ar is not a 2-ethyl-6-methyl-phenyl radical, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and its salts.

2. A compound of the formula 1 as claimed in claim 1,

where

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

Ar is one with R4, R5, R6 and R7 substituted mono- or bicyclic aromatic residue from the group of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

R6 is hydrogen,

R7 is hydrogen,

with the proviso that, in the case when Ar is not a 2-ethyl-6-methyl-phenyl radical, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and its salts.

3. A compound of the formula 1 as claimed in claim 1,
where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted monocyclic aromatic residue selected from the group of phenyl, pyridinyl, thiophenyl, furanyl and pyrrolyl,
wherein

R4 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

R5 is hydrogen, 1-4C-alkyl, halogen or fluoro-1-4C-alkyl,

R6 is hydrogen

R7 is hydrogen

and its salts.

4. A compound of the formula 1 as claimed in claim 1,
where

R1 is 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino or piperazino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-7C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted phenyl,

wherein

R4 is hydrogen or 1-4C-alkyl

R5 is hydrogen or 1-4C-alkyl

R6 is hydrogen

R7 is hydrogen

and its salts.

5. A compound of the formula 1 as claimed in claim 1,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo or morpholino radical and is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted phenyl,

wherein

R4 is hydrogen or 1-4C-alkyl

R5 is hydrogen or 1-4C-alkyl

R6 is hydrogen

R7 is hydrogen

and its salts.

6. A compound of the formula 1 as claimed in claim 1,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR31R32,

where

R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

Ar is one with R4, R5, R6 and R7 substituted phenyl,

wherein

R4 is hydrogen or 1-4C-alkyl


R5 is hydrogen or 1-4C-alkyl

R6 is hydrogen

R7 is hydrogen

and its salts.

7. A compound of the formula 1 as claimed in claim 1, characterized by the formula 1-a



where

- R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

R5 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, halogen or fluoro-1-4C alkyl

with the proviso that, in the case when R4 is not ethyl and R5 is not methyl, R1 does not have the meaning hydrogen or 1-4C-alkyl when R2 has the meaning halogen or fluoro-1-4C-alkyl, and its salts.

8. A compound of the formula 1 as claimed in claim 1, characterized by the formula 1-a as claimed in claim 7,

where


R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR31R32,

where

- R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,
 R4 is hydrogen or 1-4C-alkyl
 R5 is hydrogen or 1-4C-alkyl
 and its salts.

9. A compound of the formula 1 as claimed in claim 1, characterized by the formula 1-b



in which

- R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl or fluoro-1-4C-alkyl,
 R2 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl, amino-1-4C-alkyl, mono- or di-1-4C-alkylamino-1-4C-alkyl, the radical Res-1-4C-alkyl or the radical -CO-NR31R32,

where

Res is a imidazo, morpholino, aziridino, azetidino, pyrrolidino, pyrrolo, piperidino, piperazino or a with R30 substituted benzylamino radical and the radical Res is bonded via its nitrogen atom or one of its nitrogen atoms to the 1-4C-alkyl radical

where

R30 is 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen or hydroxy,

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino, morpholino, aziridino or azetidino group,

and its salts.

10. A compound of the formula 1 as claimed in claim 1, characterized by the formula 1-b as claimed in claim 9,

where

R1 1-4C-alkyl

R2 is carboxyl, 1-4C-alkoxycarbonyl or the radical -CO-NR31R32,

where

R31 is 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

R32 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

and its salts.

11. A medicament comprising a compound as claimed in claim 1 and/or a pharmacologically acceptable salt thereof together with customary pharmaceutical auxiliaries and/or excipients.

12. The use of a compound as claimed in claim 1 and its pharmacologically acceptable salts for the prevention and treatment of gastrointestinal disorders.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/050697

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/04 A61P1/04 A61K31/424 C07D333/38 C07D333/24
C07D277/56 C07D277/30 C07D213/87 C07D213/54 C07D215/54
C07D215/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 358 453 A (BRISTOL, JAMES A. ET AL) 9 November 1982 (1982-11-09) cited in the application claims 1,8-12,16; example -----	1,11,12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

12 May 2005

Date of mailing of the international search report

25/05/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No
PCT/EP2005/050697

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4358453	A	09-11-1982	NONE